

CLIMATE CHANGE AND AGRICULTURE RESEARCH PAPER

Statistical modelling of grapevine phenology in Portuguese wine regions: observed trends and climate change projections

H. FRAGA^{1*}, J. A. SANTOS¹, J. MOUTINHO-PEREIRA¹, C. CARLOS², J. SILVESTRE³,
J. EIRAS-DIAS³, T. MOTA⁴ AND A. C. MALHEIRO¹

¹ Centre for the Research and Technology of Agro-Environmental and Biological Sciences, Universidade de Trás-os-Montes e Alto Douro, 5000-801 Vila Real, Portugal

² Associação para o Desenvolvimento da Viticultura Duriense, Quinta da Santa Maria, 5050-106 Godim, Portugal

³ Instituto Nacional de Investigação Agrária e Veterinária, I.P., Quinta da Almoinha, 2565-191 Dois Portos, Portugal

⁴ Comissão de Viticultura da Região dos Vinhos Verdes, Estação Vitivinícola Amândio Galhano, Paçô, 4970-249 Arcos de Valdevez, Portugal

(Received 17 March 2015; revised 28 July 2015; accepted 25 August 2015)

SUMMARY

Phenological models are considered key tools for the short-term planning of viticultural activities and long-term impact assessment of climate change. In the present study, statistical phenological models were developed for budburst (BUD), flowering (FLO) and veraison (VER) of 16 grapevine varieties (autochthonous and international) from the Portuguese wine-making regions of Douro, Lisbon and Vinhos Verdes. For model calibration, monthly averages of daily minimum (Tmin), maximum (Tmax) and mean (Tmean) temperatures were selected as potential regressors by a stepwise methodology. Significant predictors included Tmin in January–February–March for BUD, Tmax in March–April for FLO, and Tmin, Tmax and Tmean in March–July for VER. Developed models showed a high degree of accuracy after validation, representing 0.71 of total variance for BUD, 0.83 for FLO and 0.78 for VER. Model errors were in most cases < 5 days, outperforming classic growing degree-day models, including models based on optimized temperature thresholds for each variety. Applied to the future scenarios RCP4.5/8.5, projections indicate earlier phenophase onset and shorter interphases for all varieties. These changes may bring significant challenges to the Portuguese wine-making sector, highlighting the need for suitable adaptation/mitigation strategies, to ensure its future sustainability.

INTRODUCTION

Grapevine development can be divided into periodic (phenological) events, which are influenced strongly by weather and climate (Jones & Davis 2000; Parker *et al.* 2011; Webb *et al.* 2012). Monitoring key phenological stages is crucial for the production of high-quality grapes and wines (Malheiro *et al.* 2013; Sadras & Moran 2013). Several morphological and physiological differences between varieties result in variances in phenological timings (Winkler *et al.* 1974). This is an additional challenge for planning management activities with potentially high economic impact for growers (Webb *et al.* 2008; Chevet *et al.* 2011), such as treatment applications (spraying) and

harvest (Tomasi *et al.* 2011; Real *et al.* 2015). Therefore, information on the development stages of different varieties, as well as the early detection of the advancement/delay of these stages, becomes increasingly important (Moriondo *et al.* 2013).

There are three main grapevine phenological stages commonly considered in the literature (Lorenz *et al.* 1995): budburst (BUD), flowering (FLO) and veraison (VER). Budburst marks the beginning of grapevine seasonal growth and resumed physiological activity, after a long period of winter dormancy; FLO is crucial for the reproductive cycle, being closely followed by the fruit set stage (Fernandez-Gonzalez *et al.* 2011) and VER initiates the ripening stage, which is tied strictly to wine grape quality attributes. The progress of these three stages summarize the main differences in varietal development throughout the growing

* To whom all correspondence should be addressed. Email: hfraga@utad.pt

season (Winkler *et al.* 1974; Fraga *et al.* 2014a), which in turn influences wine quality attributes (Jones & Davis 2000).

Climate is a major forcing factor on grapevine development and growth. Amongst the more important climatic factors, air temperature plays a leading role in governing grapevine phenology (Dalla Marta *et al.* 2010; Bonnefoy *et al.* 2013; Bonada & Sadras 2015), greatly influencing the timing of phenological stages (McIntyre *et al.* 1982; Neumann & Matzarakis 2014). Additionally, temperature largely bounds the geographical distribution of grapevines (Fraga *et al.* 2014b), also showing a deep connection to the variability in grapevine yield (Camps & Ramos 2012; Bock *et al.* 2013), wine production (Santos *et al.* 2013; Fraga *et al.* 2014c) and quality (White *et al.* 2006; de Orduna 2010).

Based on these temperature–grapevine relationships, statistical modelling may be used to predict phenology (de Cortazar-Atauri *et al.* 2009; Parker *et al.* 2011, 2013; Moriondo *et al.* 2015). Phenology models are indeed key tools for short-term planning of viticultural activities and for studying long-term climate change impacts (Caffarra & Eccel 2011). Previous studies (van Leeuwen *et al.* 2008; Nendel 2010; Parker *et al.* 2011; Fila *et al.* 2014; Neumann & Matzarakis 2014) applied conventional models based on growing degree-day (GDD) measures. Growing degree-days reflect the amount of accumulated heat above the base temperature of 10 °C (Winkler *et al.* 1974) in each growth stage. However, recent research hints at the limitations of these GDD models, such as the need for site-specific calibration and uncertainties in the temperature threshold (de Cortazar-Atauri *et al.* 2009; Caffarra & Eccel 2010, 2011; Parker *et al.* 2011; Zapata *et al.* 2015). Most of these models have been applied to regions where climatic conditions are very different from the Mediterranean-like conditions prevailing in Portugal. Furthermore, it has been shown that temperatures over critical intervals may strongly influence grapevine development, which are not adequately captured by the accumulation of daily temperatures (Malheiro *et al.* 2013).

In the climate change context, several studies have been carried out to investigate its impacts on viticulture. Climate change has the potential to modify this crop markedly, providing additional challenges for wine growers (Jones *et al.* 2005b; Moriondo & Bindi 2007; Malheiro *et al.* 2010; Hannah *et al.* 2013). Regarding phenology, most studies project earlier phenological timings and shorter intervals under

future warmer climates (Jones *et al.* 2005a; Webb *et al.* 2007, 2011; Tomasi *et al.* 2011). However, climate change impacts on viticulture are known to be more tied to regional/local climatic conditions than to large-scale changes (Anderson *et al.* 2012; Fraga *et al.* 2015b). Therefore, regional assessments of climate change impacts on vine phenology need to be undertaken. No previous research has been focused on the impacts of climate change on grapevine phenology in the Portuguese wine regions. Hence, a deeper understanding of specific varietal differences in phenology is critical to select the most suitable varieties under future climatic conditions (Duchene *et al.* 2010).

The objectives of the present study are threefold: (1) to develop a modelling approach that captures the variability of the main grapevine (mostly native) varieties grown in Portugal; (2) to validate the models using different varieties from different Portuguese regions and compare their accuracy with the more conventional GDD models; and (3) to examine the impacts of climate change on regional grapevine phenology.

MATERIALS AND METHODS

Phenology and climatic data

Phenological data were collected from three different wine-making regions (Fig. 1): Douro (Northeastern Portugal), Vinho Verdes (Minho, Northwestern Portugal) and Lisbon (Centralwestern Portugal). These three regions, with centuries-old wine-making traditions, jointly have an important share of the national wine-making economy. The Douro Demarcated Region (including the Denominations of Origin Douro and Porto) is currently the main viticultural region in the country in terms of wine production and vineyard area (150 million litres and 45 000 ha), followed by Lisbon (90 million litres and 25 000 ha) and the Demarcated Region of Vinhos Verdes (80 million litres and 31 000 ha) (IVV 2013). These world-renowned regions are best known for their Port (Douro) and Vinho Verde wines, of which roughly half of the annual production is exported (IVV 2013).

Table 1 describes the 16 grapevine varieties selected for the current study, as well as the corresponding time periods with phenological data. For the Vinhos Verdes region, phenological timings were recorded for the Loureiro (white) variety in the period 2003–2012, from a vineyard located at ‘Arcos de Valdevez’ (41°48′57″ N, 8°24′35″W, 70 m a.s.l.). For Douro, phenological data were collected for the Touriga-Franca (red)




Fig. 1. Location of vineyard sites where data was collected. The delimitations indicate the borders of the winemaking regions in Portugal, as defined by the Portuguese 'Instituto do Vinho e da Vinha'.

variety in the period 2001–2011, at 'Ervedosa do Douro' (41°10'31"N, 7°32'58"W, 180 m a.s.l.). For Lisbon, data for seven white (Alvarinho, Encruzado, Fernão-Pires, Rabigato, Riesling, Trincadeira-das-Pratas and Viosinho) and seven red (Borraçal, Castelão, Grenache, Jaen, Merlot, Pinot and Tinta-Francisca) varieties were collected from an experimental vineyard located at Dois Portos (DP) (39°02'35"N, 9°10'55"W, 80 m a.s.l.) over the combined period 1991–2011. The timings of BUD, FLO and VER were recorded through field observations following the grapevine descriptors recommended by the Organisation Internationale de la Vigne et du Vin (OIV 2009). The mean calendar day of year (DOY; 1 corresponds to 1 January of a given year) of each phenophase for each variety are also presented in Table 1.

The three target regions are characterized by Mediterranean-like climates, with warm dry

summers and mild wet winters (Peel *et al.* 2007). According to the 1981–2010 climatic norms provided by the Portuguese meteorological office (Instituto Português do Mar e da Atmosfera (IPMA); <http://www.ipma.pt>), the Vinhos Verdes region typically shows annual mean temperatures of about 15 °C, Douro of 13 °C and Lisbon of 17 °C. For the subsequent analysis, historical records of daily minimum (Tmin), maximum (Tmax) and mean (Tmean) air temperatures, covering the study period and representative of the vineyard sites, were obtained. For Vinhos Verdes, data were obtained from the Sistema Nacional de Informação de Recursos Hídricos (SNIRH) database (<http://snirh.apambiente.pt>) for the Ponte da Barca weather station, PB (SNIRH code 03G/02C), only 2 km from the vineyard. For the Douro site, due to the lack of local data, climate records were obtained from IPMA for the Vila Real weather station, VR (IPMA code 80567), located about 20 km north-west of the vineyard. For the Lisbon site, data were obtained from the weather station installed in the vineyard, situated at DP.

Phenological models

Each phenophase was modelled for white and red varieties separately, as important differences in the timings of these two types of grape can occur. Although differences in phenological timings may also occur between varieties within these two sub-groups (de Cortazar-Atauri *et al.* 2009b), it was assumed that these differences could be captured by the present models. For this purpose, the varieties with the longest time series of phenological data (Table 1) were selected, i.e. Fernão-Pires (white) and Castelão (red). The full raw time series of the yearly DOY of BUD, FLO and VER were used as dependent variables in linear regression models (least squares estimation).

In order to establish statistical relationships between temperature and phenology, monthly averages of daily mean (Tmean), minimum (Tmin) and maximum (Tmax) air temperatures from each site were used as independent variables in the models (predictors). Due to the large number of potential predictors, a stepwise multivariate regression was then undertaken for variable selection (Wilks 2011). The stepwise criterion for forward inclusion and backward removal correspond to $P < 0.05$ and $P < 0.10$, respectively. The initial set of potential predictors comprised not only monthly means of Tmin, Tmax and Tmean preceding each phenophase but also multi-month mean temperatures. The

Table 1. Mean day of year (DOY, calendar day of year where 1 corresponds to 1 Jan of a given year) of the BUD (Budburst), FLO (Flowering) and VER (Veraison) for varieties used in the study. Time interval of each series is also shown

Whites varieties	Years	Region	Mean DOY		
			BUD	FLO	VER
Alvarinho	1995–2011	Lisbon	77	142	221
Encruzado	1995–2005	Lisbon	77	141	223
Fernão-Pires	1990–2011	Lisbon	75	142	215
Loureiro	2003–2012	Vinhos Verdes	85	150	222
Rabigato	1995–2005	Lisbon	72	143	223
Riesling	1990–2000	Lisbon	79	142	218
Trincadeira-das-Pratas	1990–2002	Lisbon	69	141	212
Viosinho	1995–2005	Lisbon	76	139	217
Red varieties					
Borraçal	1995–2005	Lisbon	74	138	226
Grenache	1990–2000	Lisbon	75	142	229
Castelão	1990–2011	Lisbon	72	139	217
Jaen	1995–2011	Lisbon	78	143	209
Merlot	1994–2005	Lisbon	75	141	215
Pinot	1993–2003	Lisbon	71	136	211
Tinta-Francisca	1995–2005	Lisbon	78	144	213
Touriga-Franca	2001–2011	Douro	85	144	207

Table 2. List of potential regressors used for model development

Phenophase	Potential regressors
Budburst	Tmin _{Jan} ; Tmin _{Feb} ; Tmin _{Mar} ; Tmin _{Jan-Feb} ; Tmin _{Feb-Mar} ; Tmin _{Jan-Mar} ; Tmax _{Jan} ; Tmax _{Feb} ; Tmax _{Mar} ; Tmax _{Jan-Feb} ; Tmax _{Feb-Mar} ; Tmax _{Jan-Mar} ; Tmean _{Jan} ; Tmean _{Feb} ; Tmean _{Mar} ; Tmean _{Jan-Feb} ; Tmean _{Feb-Mar} ; Tmean _{Jan-Mar}
Flowering	Tmin _{Mar} ; Tmin _{Apr} ; Tmin _{May} ; Tmin _{Mar-Apr} ; Tmin _{Apr-May} ; Tmin _{Mar-May} ; Tmax _{Mar} ; Tmax _{Apr} ; Tmax _{May} ; Tmax _{Mar-Apr} ; Tmax _{Apr-May} ; Tmax _{Mar-May} ; Tmean _{Mar} ; Tmean _{Apr} ; Tmean _{May} ; Tmean _{Mar-Apr} ; Tmean _{Apr-May} ; Tmean _{Mar-May}
Veraison	Tmin _{Mar} ; Tmin _{Apr} ; Tmin _{May} ; Tmin _{Jun} ; Tmin _{Jul} ; Tmin _{Mar-Apr} ; Tmin _{Apr-May} ; Tmin _{May-Jun} ; Tmin _{Jun-Jul} ; Tmin _{Mar-Jul} ; Tmax _{Mar} ; Tmax _{Apr} ; Tmax _{May} ; Tmax _{Jun} ; Tmax _{Jul} ; Tmax _{Mar-Apr} ; Tmax _{Apr-May} ; Tmax _{May-Jun} ; Tmax _{Jun-Jul} ; Tmax _{Mar-Jul} ; Tmean _{Mar} ; Tmean _{Apr} ; Tmean _{May} ; Tmean _{Jun} ; Tmean _{Jul} ; Tmean _{Mar-Apr} ; Tmean _{Apr-May} ; Tmean _{May-Jun} ; Tmean _{Jun-Jul} ; Tmean _{Mar-Jul}

full list of potential predictors for each phenophase is shown in Table 2. Based on the predictors selected by the stepwise methodology, a linear model was then fitted for each of the phenological stages. A leave-one-out cross-validation scheme was applied to account for model overfitting. Model residuals were tested using both the Durbin–Watson test and the corresponding autocorrelation function, showing no serial correlation between them (independently distributed).

Model performance

For the assessment of model accuracy, the determination coefficient (R_{CV}^2 ; after a leave-one-out cross-

validation) and the root-mean-squared error (RMSE) were estimated. An out-of-sample validation was also performed by applying the developed models to other grapevine varieties grown in the Lisbon region and in the other two regions (Vinhos Verdes and Douro), allowing an inter-regional assessment of the model accuracy. Subsequently, these models were compared against a standard GDD model to test their added-value. The GDD model was computed for both Fernão-Pires and Castelão varieties using the available climatic data from Lisbon. The mean GDD for BUD, FLO and VER stages were computed for 1990–2011, using the standard 10 °C base temperature (Winkler *et al.* 1974) and an estimated base

temperature by applying the minimizing the standard deviation method (Zapata *et al.* 2015). For Fernão-Pires the computed base temperature, using this method, was 9.2 °C, while for Castelão it was 8.9 °C. The GDD values, using these base temperatures (10 °C and 9.2/8.9 °C), were then used to determine the yearly DOY of each phenophase. These models were then compared to the newly developed models using R_{cv}^2 and RMSE.

Future projections

Following the phenological model calibration and validation, an analysis of climate change impacts on the phenological timings and intervals was performed. For this purpose, the developed phenological models are then applied to future projections of the mentioned predictors. Four pairs of Global Climate Model/Regional Climate Model simulations (Table 3) were retrieved from the EURO-CORDEX project (Jacob *et al.* 2014). All simulations cover the Euro-Atlantic domain at a spatial resolution of $0.11^\circ \times 0.11^\circ$ (~12.5 km). These simulations integrate the Coupled Model Intercomparison Project (CMIP5) and are forced by two Representative Concentration Pathways (RCP) scenarios – RCP4.5 and RCP8.5 for 2006–2100. All models were previously subjected to calibration and bias correction using the Era-Interim reanalysis (Jacob *et al.* 2014; Kotlarski *et al.* 2014). As the number of simulations was relatively large, only the ensemble-mean of each scenario for 2020–2090 was used herein to isolate the climate change signal and to assess regional-scale uncertainties.

RESULTS

Present climatic conditions

The three selected wine regions, PB–Vinhos Verdes, VR–Douro and DP–Lisbon, revealed similar temperature seasonality (Fig. 2(a) and (c) and (e)). Ponte da Barca–Vinhos Verdes presented average annual minima of 9.6 °C, maxima of 20.7 °C and mean of 14.7 °C. Vila Real–Douro showed much lower temperatures, with annual minima of 8.8 °C, maxima of 18.9 °C and mean of 13.8 °C. Dois Portos–Lisbon presented temperature values very close to PB–Vinhos Verdes in terms of annual maxima with 20.7 °C, but with a higher minima of 11.1 °C and mean of 15.9 °C. Temperatures are usually highest in August and lowest in January/December in all regions. Vila Real–Douro

presented the highest thermal amplitudes (Fig. 2(c)). Dois Portos–Lisbon presented the mildest temperatures during winter (Fig. 2(e)), while PB–Vinhos Verdes exhibited intermediate temperatures. It should be noted that for DP–Lisbon, monthly mean temperatures during the study period (1990–2011) were always above 10 °C, which is commonly considered a temperature threshold for grapevine development (Winkler *et al.* 1974). Regarding the annual mean temperature trends (Fig. 2(b) and (d) and (f)), only Lisbon showed statistically significant trends at $P < 0.05$ (Fig. 2(f)), with values for Tmean and Tmax of 0.04 and 0.03 °C/year, respectively.

Phenological models

As mentioned above, the phenological series of one white (Fernão-Pires) and one red (Castelão) variety were selected for a first model calibration. Their phenological time series are shown in Fig. 3. No significant trends were found in the phenological stages of either variety, according to the non-parametric Mann–Kendall test. Regarding the mean DOY of the two grapevine varieties (Table 1), Fernão-Pires tended to reach both BUD and FLO 2–3 days later than Castelão, while Castelão tended to reach VER 2 days later than Fernão-Pires (Table 1). Fernão-Pires showed a lower variability in BUD than Castelão, with interquartile-ranges (IQR) of 10 and 12 days, respectively (Fig. 3). Both varieties showed the same 7-day IQR for FLO. For VER, Castelão and Fernão-Pires show higher inter-annual variability (12 and 10 days of IQR, respectively). The inter-annual variability of these last two events were particularly evident in the year 1997, with an anomalously warm spring. On the other hand, other warmer years, such as 2003, had little influence on the studied phenology, since heat waves occurred mainly after VER (August). Regarding the BUD–FLO interphase length, Fernão-Pires showed a mean duration of 67 days (IQR = 11 days), the same as Castelão (IQR = 10 days). For FLO–VER, Castelão showed higher mean interphase duration than Fernão-Pires (78 and 73 days, respectively) and lower IQR (7 and 8 days, respectively).

Following the stepwise process, the most significant predictors were then isolated for modelling the individual phenophases for each variety. For BUD, the identified predictors were $T_{min_{Feb-Mar}}$ for the white model and $T_{min_{Jan-Mar}}$ for the red model. For FLO, the same predictor was identified in both models (white and red): $T_{max_{Mar-Apr}}$. For VER, the stepwise

Table 3. Selected global/regional climate model chains over the period of 2020–2089. All model simulations belong to the Euro-Cordex project (EUR-11 ensemble) and are available at the original grid of 0.11° latitude × 0.11° longitude (spatial resolution of c. 12.5 km)

GCM	Institution	RCM	Institution
CNRM– CERFACS	Centre National de Recherches Météorologiques	SMHI–RCA4	Swedish Meteorological and Hydrological Institute
EC–EARTH	Royal Netherlands Meteorological Institute	DMI– HIRHAM5	Danish Meteorological Institute
MOHC– HadGEM2	Met Office Hadley Centre	SMHI–RCA4	Swedish Meteorological and Hydrological Institute




Fig. 2. Left panel: Monthly averages of daily minimum (Tmin), mean (Tmean) and maximum (Tmax) air temperatures registered at (a) Vinhos Verdes, (c) Douro and (e) Lisbon. Right panel: Chronogram of the annual averaged daily minimum (Tmin), mean (Tmean) and maximum (Tmax) air temperatures registered at (b) Vinhos Verdes, (d) Douro and (f) Lisbon.




Fig. 3. (a) *Left panels:* Chronograms of the calendar day of year (DOY) of budburst (BUD), flowering (FLO) and veraison (VER) for Fernão-Pires and over the period of 1990–2011, along with the respective linear trends (LT). No statistically significant LT is found; *Middle panels:* Histograms of the frequencies of occurrence of DOY for each phenophase. The corresponding mean (+), median (vertical bar), 25th and 75th percentiles (box limits), 9th and 91st percentiles (whiskers) and outliers (circles) are also shown above each histogram; *Right panels:* the same as on left panel, but for the respective interphase durations. (b) As in (a), but for Castelão.

process identified the largest differences between varieties as well as more than one significant regressor per model. For the white model, the first regressor was $T_{\min_{\text{Mar-Jul}}}$ and the second regressor $T_{\max_{\text{Mar-Apr}}}$.

For the red model, the first regressor was $T_{\max_{\text{Jun-Jul}}}$ and the second $T_{\text{mean}_{\text{Mar-Apr}}}$. The corresponding linear models were significant at a $P < 0.001$ and are represented in Eqns (1) and (3) for the white varieties

and in Eqns (4) and (6) for the red varieties:

$$\text{BUD}_{\text{White}} = k + \alpha \times \text{Tmin}_{\text{Feb-Mar}} \quad (1)$$

$$\text{FLO}_{\text{White}} = k + \alpha \times \text{Tmax}_{\text{Mar-Apr}} \quad (2)$$

$$\text{VER}_{\text{White}} = k + \alpha \times \text{Tmin}_{\text{Mar-Jul}} + \beta \times \text{Tmax}_{\text{Mar-Apr}} \quad (3)$$

$$\text{BUD}_{\text{Red}} = k + \alpha \times \text{Tmin}_{\text{Jan-Mar}} \quad (4)$$

$$\text{FLO}_{\text{Red}} = k + \alpha \times \text{Tmax}_{\text{Mar-Apr}} \quad (5)$$

$$\text{VER}_{\text{Red}} = k + \alpha \times \text{Tmax}_{\text{Jun-Jul}} + \beta \times \text{Tmean}_{\text{Mar-Apr}} \quad (6)$$

where k corresponds to the intercept (regression constant), α to the first regressor coefficient and β to the second regressor coefficient, if one exists.

Model validation

Following the identification of the most significant predictors and model development, model accuracy was determined by the previously mentioned metrics (R_{cv}^2 and RMSE). Figure 4 depicts observed *v.* modelled phenological timings (DOY of each phenological event). Overall, the models performed very well, with high R_{cv}^2 and relatively low RMSE for all phenological stages. The FLO and VER models performed better than BUD models in depicting the observed DOY for both varieties, with lower R_{cv}^2 but similar accuracy (RMSE). The BUD models showed an R_{cv}^2 of 0.63 for Fernão-Pires and of 0.69 for Castelão. Nonetheless, the RMSE for these models was low (4–5 days). The FLO models for both varieties showed higher accuracy, with an R_{cv}^2 of 0.79 and RMSE lower than 4 days. For VER, the models showed an R_{cv}^2 of 0.77 for Fernão-Pires (RMSE = 5 days) and of 0.88 for Castelão (RMSE < 3 days). Comparing the results of the newly developed models against a standard GDD model (Table 4), the former tended to outperform the latter, showing higher R_{cv}^2 and lower RMSE, for both varieties and all phenological stages.

For validation purposes, the same models were then applied to other varieties growing not only in the Lisbon wine region (Tables 5 and 6), but also in Vinhos Verdes and Douro (Table 7). In general, for Lisbon, the model performance was very high for all

phenological stages and varieties. White varieties showed, on average, higher performance in BUD and VER and red varieties in FLO. Nevertheless, the performance of the BUD model is slightly lower than of the FLO and VER models. Average R_{cv}^2 for BUD white/red models was 0.70/0.85 and RMSE of 4/3 days. The FLO models showed the best performances, with a varietal average R_{cv}^2 of 0.87/0.89 (white/red varieties) and RMSE of 2–5 days. The VER models show average white/red R_{cv}^2 of 0.86/0.82 and average RMSE of 5/4 days.

When applying the same models to other varieties growing in other regions (Table 7), model performance generally decreased (R_{cv}^2), but the accuracy (RMSE) remained almost unchanged, with the exception of the BUD model for Loureiro, which exceeded 6 days. Generally, BUD models (as previously) showed the lowest R_{cv}^2 of the three phenophase models, with 0.32 for Loureiro (white) and 0.38 for Touriga-Franca. For FLO, the model showed a relatively high performance for Touriga-Franca ($R_{cv}^2 = 0.74$), but much lower for Loureiro ($R_{cv}^2 = 0.50$). For the VER model, the performance was similar in both regions/varieties, with an R_{cv}^2 of 0.59 for Loureiro and of 0.51 for Touriga-Franca. Despite the relatively low performance of the BUD model when applied to other regions/varieties, the FLO and VER models explained more than 50% of the represented variance in the phenological timings, which was a noteworthy outcome.

Temperature projections

Analysis of future projections for air temperature in mainland Portugal revealed a strong warming trend (Fig. 5). Therefore, for the sake of succinctness, only the temperature projections for the main predictors of the phenological models in the Lisbon wine region will be analysed, as similar warming trends are expected for Vinhos Verdes and Douro (cf. Fig. 5). Temperature projections for experiments under the RCP4.5 and RCP8.5 scenarios reveal enhanced upward/warming trends with respect to recent-past trends (Fig. 6): RCP4.5 shows a significant increase of 0.01–0.02 °C/year while the RCP shows increases of 0.03–0.05 °C/year for the selected predictors. Therefore, based on these projections, the most severe emission scenario (RCP8.5) shows the highest warming trends, as expected.

Phenological projections

An analysis of the phenological projections was performed taking into account the period of 2040–




Fig. 4. *Left panel:* Scatterplots of modelled v. observed calendar day of year (DOY) of budburst (BUD), flowering (FLO) and veraison (VER) for Fernão-Pires over the period 1990–2011 (22 years). The corresponding linear regression line is also plotted. *Right panel:* the same as on left panel, but for Castelhão.

Table 4. *Comparison between the growing degree-day model (GDD), minimizing standard-deviation (SD) model, along with the calculated base temperature, and regression models, for each phenophase of Fernão-Pires and Castelhão. Accuracy parameters for all models are shown: R_{cv}^2 and RMSE*

	GDD model (10 °C)			Minimizing SD model			Regression models	
	Mean GDD	R_{cv}^2	RMSE	Base Temp.	R_{cv}^2	RMSE	R_{cv}^2	RMSE
Budburst								
Fernão-Pires	223	0.40	5.69	9.2	0.22	7.38	0.63	4.48
Castelhão	208	0.48	6.01	8.9	0.31	7.06	0.69	4.63
Flowering								
Fernão-Pires	671	0.67	4.85	9.2	0.73	4.69	0.79	3.85
Castelhão	645	0.62	5.19	8.9	0.70	4.30	0.79	3.90
Veraison								
Fernão-Pires	1551	0.31	8.62	9.2	0.35	5.49	0.77	5.00
Castelhão	1573	0.53	5.75	8.9	0.57	8.26	0.88	2.89

Table 5. White grapevine phenology model for six varieties grown in the Lisbon winemaking region. The R_{cv}^2 , root-mean-square error (RMSE, in days), k – intercept, α – first regressor coefficient and β – second regression coefficient are also shown. The independent variables for modelling each phenophase–variety pair are shown in Eqns (1) and (3)

Budburst	R_{cv}^2	RMSE	k	α	β
Alvarinho	0.65	4.31	123.68	–6.10	
Encruzado	0.84	2.77	123.39	–5.88	
Rabigato	0.75	4.03	123.31	–6.51	
Riesling	0.65	4.03	117.22	–5.09	
Trincadeira-das-Pratas	0.68	3.98	111.42	–5.42	
Viosinho	0.68	4.18	120.10	–5.69	
Flowering					
Alvarinho	0.87	2.98	291.44	–7.89	
Encruzado	0.91	2.90	293.56	–8.07	
Rabigato	0.90	3.01	298.45	–8.25	
Riesling	0.84	4.89	305.29	–8.69	
Trincadeira-das-Pratas	0.80	4.27	267.70	–6.79	
Viosinho	0.85	3.57	285.96	–7.80	
Veraison					
Alvarinho	0.75	5.09	457.65	–14.27	–3.54
Encruzado	0.75	5.29	401.28	–3.84	–7.01
Rabigato	0.75	5.64	458.63	–10.18	–6.05
Riesling	0.89	3.46	407.18	–6.08	–6.21
Trincadeira-das-Pratas	0.83	4.35	439.08	–12.42	–4.30
Viosinho	0.85	4.50	456.85	–9.028	–6.99

2070, in comparison to the current (1991–2011) phenological observations. For all varieties studied, the mean DOY of all phenological timings and intervals showed advances, i.e. earlier timings (Fig. 7). The advancement was more pronounced for scenario RCP8.5 than for RCP4.5 and for the VER stage. Both white and red varieties seemed to be equally affected in the FLO and VER stage, but differently in the BUD stage. For BUD, the advance was slightly more pronounced in the red varieties: 1–5 v. 1–3 days in the white varieties (Fig. 7). The FLO model showed 2–6 days advancement, while VER showed the largest changes, with an advance of 6–14 days depending on future scenario. Concerning specific varietal changes, Fernão-Pires and Grenache showed the largest advances, while Encruzado and Jaen showed the smallest advances of all varieties studied. Future changes hint at significant shortenings of the phenological timings and intervals, while the enhanced warming conditions for grapevine growth will contribute to longer growing seasons. A reduction of 1–2 days could be expected in BUD–FLO for red varieties and 1–3 days for white varieties. This red/white difference can be explained by the more pronounced

advancement of the red varieties at the BUD timings. For the FLO–VER interval, a 4–8-day shortening is projected under the RCP4.5–RCP8.5 scenarios.

DISCUSSION

The present study focused on model development for both short-term phenological prediction and long-term climate change impact assessment: it investigates comprehensive changes in future climate under the emission scenarios RCP4.5 and RCP8.5 and their impact on grapevine phenology. Several previous studies have highlighted the impact of climate change on phenology, such as earlier phenophases and shorter intervals (Webb *et al.* 2007; Hall & Jones 2009; Duchene *et al.* 2010). However, no studies focusing on climate change impact on the phenology of Portuguese grapevine varieties has been undertaken so far. Furthermore, no previous phenological model has been simultaneously applied to several varieties of different wine-making regions in Portugal.

Using the longest available phenological records in Portugal, a modelling approach was developed accounting for a broad number of varieties and

Table 6. Red grapevine phenology model for six varieties grown in the Lisbon winemaking region. The R_{cv}^2 , root-mean-square error (RMSE, in days), k – intercept, α – first regressor coefficient and β – second regression coefficient are also shown. The independent variables for modelling each phenophase–variety pair are shown in Eqns (4) and (6)

Budburst	R_{cv}^2	RMSE	k	α	β
Borraçal	0.93	2.25	143.20	−9.48	
Grenache	0.91	2.25	125.83	−7.44	
Jaen	0.70	4.35	136.47	−8.09	
Merlot	0.80	3.65	137.69	−8.69	
Pinot	0.84	2.94	129.85	−8.05	
Tinta-Francisca	0.91	2.62	147.38	−9.53	
Flowering					
Borraçal	0.92	3.03	309.20	−9.05	
Grenache	0.83	4.77	294.55	−8.16	
Jaen	0.90	2.68	300.48	−8.30	
Merlot	0.89	3.26	309.54	−8.92	
Pinot	0.84	4.55	302.30	−8.84	
Tinta-Francisca	0.93	2.68	307.12	−8.61	
Veraison					
Borraçal	0.89	3.38	488.75	−3.24	−12.81
Grenache	0.78	3.79	586.18	−7.23	−12.78
Jaen	0.80	3.56	388.39	−1.57	−9.87
Merlot	0.74	5.18	554.83	−5.62	−13.94
Pinot	0.80	4.04	419.65	−2.75	−9.98
Tinta-Francisca	0.91	2.36	396.83	−1.89	−9.60

Table 7. Phenology models applied to varieties in other viticultural regions: Loureiro from Vinhos Verdes region (white variety) and Touriga-Franca from Douro region (red variety). The R_{cv}^2 , root mean square error (RMSE, in days), k – constant, α – first regressor coefficient and β – second regression coefficient are also shown

Budburst	R_{cv}^2	RMSE	k	α	β
Loureiro	0.32	6.65	108.52	−4.76	
Touriga-Franca	0.38	3.59	100.15	−3.64	
Flowering					
Loureiro	0.50	4.71	248.18	−5.34	
Touriga-Franca	0.74	2.91	229.63	−5.25	
Veraison					
Loureiro	0.59	4.43	350.71	−4.50	−4.37
Touriga-Franca	0.51	4.73	318.28	−1.09	−7.19

regions. The model was successful in predicting the phenological timings of BUD, FLO and VER. Overall, for the 16 varieties studied, the average R_{cv}^2 for BUD was 0.71, for FLO 0.83 and for VER 0.78. For all phenological stages, the average RMSE was < 5 days in most cases, but for one variety (Loureiro) at BUD. These outcomes are a noteworthy progress, being also comparable, in terms of model accuracy, to other studies in

other European wine-making regions (Bindi *et al.* 1997; de Cortazar-Atauri *et al.* 2009; Caffarra & Eccel 2010; Parker *et al.* 2011, 2013).

The models developed out-performed the classic GDD models. Although the GDD models are currently standard in the wine-making industry (Winkler *et al.* 1974; Fraga *et al.* 2015a; Zapata *et al.* 2015), several limitations to their applicability have been pointed




Fig. 5. Ensemble mean air temperature trends (annual mean temperature; °C/year) calculated using the EURO-CORDEX simulations over the period of 2020–2090 and under the RCP 4.5 (*left panel*) and RCP 8.5 (*right panel*). Colour online.

out. Some studies highlight that the GDD calculation is site-specific and cannot be extrapolated to other regions (de Cortazar-Atauri *et al.* 2009; Caffarra & Eccel 2010). Other findings demonstrate that the 10 °C base temperature in GDD models may not represent a suitable threshold for all varieties and wine-grape growing regions (Parker *et al.* 2011; Zapata *et al.* 2015). The newly developed models also outperformed the GDD models based on optimized temperature thresholds. These results, obtained for three main wine regions in Portugal, suggest that phenology exhibits a stronger relationship with temperature averages in certain specific periods than with the summations. In fact, key physiological and biochemical processes, such as the remobilization of stored starch and supply of sugar from BUD to FLO, from the permanent structure of the vine to shoot and leaf growth, are initiated by optimum temperatures, promoting expansion and differentiation (Keller 2010b).

The key periods in which phenological development is linearly tied to air temperature are herein identified, confirming the existence of significant links between grapevine phenological timing and temperature in the preceding months, in agreement with previous studies (Malheiro *et al.* 2013). Minimum temperatures from January/February to March have been shown to have a strong influence on BUD timings. Nonetheless, this stage proved to be more difficult to model, which may

be due to the influence of viticultural practices, such as pruning date (Martin & Dunn 2000) or insufficient winter chill (Webb *et al.* 2007). Maximum temperatures from March to April show a linear relationship with FLO, in agreement with other studies (Bock *et al.* 2011; Tomasi *et al.* 2011). The VER timings and associated predictors show the highest variability of all phenophases. In addition, the main climatic drivers for this phenophase differ for white and red varieties. For white varieties, the minimum temperatures from March to July and the maximum temperatures from March to April are the strongest climatic drivers. For red varieties, the main driving factors are the maximum temperature from June to July and the mean temperature from March to April. Further, the second regressor (March–April temperatures) coincides with the FLO main climatic driver, highlighting a link between these two phenophases (FLO and VER). In summary, strong links between BUD and winter temperatures, FLO and spring temperatures and VER with spring–summer temperatures were found.

The models showed lower performance when applied to other grapevine varieties growing in other regions in Portugal (Vinhos Verdes and Douro). This result suggests that models need to be locally calibrated in order to optimize their performance. However, lower accuracy in these two regions can




Fig. 6. Chronogram of historical observations (1990–2011) in the Lisbon site and corresponding future projections (2020–2090 under RCP4.5 and RCP 8.5) from the EUR-11 4-member ensemble (cf. Table 3) and for: (a) February–March average daily minimum temperature ($T_{\min_{\text{Feb-Mar}}}$); (b) March–April average daily maximum temperature ($T_{\max_{\text{Mar-Apr}}}$), (c) March–July average daily minimum temperature ($T_{\min_{\text{Mar-Jul}}}$), (d) January–March average daily minimum temperature ($T_{\min_{\text{Jan-Mar}}}$), (e) June–July average daily maximum temperature ($T_{\max_{\text{Jun-Jul}}}$) and (f) March–April average daily mean temperature ($T_{\text{mean}_{\text{Apr-Mar}}}$).

be partially attributed to the location of the corresponding weather stations (data used) outside the vineyard, which was reported to have a strong impact on modelling results (Olsson & Jönsson 2015). In Douro, the weather station is 20 km away from the vineyard site, thus being a likely source of errors.

Climate change projections applied to the selected models indicate that future warming will lead to earlier phenological timings and intervals. These outcomes suggest that the VER will have the strongest change, occurring about 2 weeks earlier. The earlier timings of these three phenophases can together reduce the grapevine development cycle considerably. These results are in agreement with recent studies worldwide, such as Duchene & Schneider (2005) for France, Jones (2005) for the USA, Webb

et al. (2007) and Petrie & Sadras (2008) for Australia and Bock *et al.* (2011) for Germany. These shifts towards earlier phenophase onsets can potentially result in changes to the currently established wine characteristics and typicity. In effect, expected warming may result in unbalanced wines, with high alcohol content and excessively low acidity, altered colour and aroma (Jones *et al.* 2004a; Malheiro *et al.* 2013). The potential effects of climate change are not exclusively limited to the reported warming trend. Other consequences projected for the Mediterranean region include higher water stress (Moriondo & Bindi 2007; Fraga *et al.* 2015b) and increased frequency of climate extremes (Santos & Corte-Real 2006; Santos *et al.* 2007; Keller 2010a), including in Portugal (Costa *et al.* 2012; Andrade *et al.* 2014). Extreme weather events, especially winter/spring weather events such as hail and frost, are




Fig. 7. *Left panel:* Differences (Future – Present: 2040–2070 minus 1990–2011) in the number of days required to reach (a) budburst (BUD), (b) flowering (FLO) and (c) veraison (VER) for the white varieties (Fernão-Pires, Alvarinho, Encruzado, Rabigato, Trincadeira-das-Pratas and Viosinho). *Right panel:* The same as on the left panel, but for the red varieties (Castelão, Borraçal, Grenache, Jaen, Merlot, Pinot and Tinta-Francisca).

projected to inflict important yield losses in several crops (IPCC 2012). Given their unpredictable nature, they may represent an additional challenge for grapevines in future climates. Nonetheless, some positive impacts are also projected, given that the overall length of the growing season is expected to increase, triggered by enhanced warming from spring to autumn.

CONCLUSIONS

In the present study, phenological models are calibrated for Portuguese grapevine varieties, out-performing standard GDD models in predicting grapevine phenology. These models provide robust results, with relatively high accuracy, enabling their

direct application by the winemaking sector. Applied to climate change scenarios, projections of the main grapevine phenological stages are achieved. These projections suggest earlier onsets for all phenological stages, as well as a shortening of the phenophase intervals, but with overall longer growing seasons. Hence, although earlier phenological stages may bring detrimental impacts for growers, some benefits may be acquired from the longer favourable periods for grapevine growth, associated with a later onset of autumn cold weather. Therefore, climate change may bring both negative and positive impacts for grapevine development, influencing its entire vegetative cycle. These findings may help delineating suitable adaptation strategies, such as a more critical varietal selection or adoption of innovative viticultural practices, which may potentiate the future sustainability of the national wine industry.

This study was supported by national funds by FCT – Portuguese Foundation for Science and Technology, under the project UID/AGR/04033/2013. This work was also supported by the project ‘ModelVitiDouro’ – PA 53774”, funded by the Agricultural and Rural Development Fund (EAFRD) and the Portuguese Government by Measure 4-1 – Cooperation for Innovation PRODER programme – Rural Development Programme. The authors thank the Estação de Avisos do Douro (Direcção Regional de Agricultura e Pescas do Norte) (Peso da Régua, Portugal) for providing the Douro’s phenological data.

REFERENCES

- ANDERSON, J. D., JONES, G. V., TAIT, A., HALL, A. & TROUGHT, M. C. T. (2012). Analysis of viticulture region climate structure and suitability in New Zealand. *Journal International des Sciences de la Vigne et du Vin* **46**, 149–165.
- ANDRADE, C., FRAGA, H. & SANTOS, J. A. (2014). Climate change multi-model projections for temperature extremes in Portugal. *Atmospheric Science Letters* **15**, 149–156.
- BINDI, M., MIGLIETTA, F., GOZZINI, B., ORLANDINI, S. & SEGHI, L. (1997). A simple model for simulation of growth and development in grapevine (*Vitis vinifera* L.). 2. Model validation. *Vitis* **36**, 73–76.
- BOCK, A., SPARKS, T., ESTRELLA, N. & MENZEL, A. (2011). Changes in the phenology and composition of wine from Franconia, Germany. *Climate Research* **50**, 69–81.
- BOCK, A., SPARKS, T. H., ESTRELLA, N. & MENZEL, A. (2013). Climate-induced changes in grapevine yield and must sugar content in Franconia (Germany) between 1805 and 2010. *Plos ONE* **8**, 10. DOI: 10.1371/journal.pone.0069015.
- BONADA, M. & SADRAS, V. O. (2015). Review: critical appraisal of methods to investigate the effect of temperature on grapevine berry composition. *Australian Journal of Grape and Wine Research* **21**, 1–17.
- BONNEFOY, C., QUENOL, H., BONNARDOT, V., BARBEAU, G., MADELIN, M., PLANCHON, O. & NEETHLING, E. (2013). Temporal and spatial analyses of temperature in a French wine-producing area: the Loire Valley. *International Journal of Climatology* **33**, 1849–1862.
- CAFFARRA, A. & ECCEL, E. (2010). Increasing the robustness of phenological models for *Vitis vinifera* cv. Chardonnay. *International Journal of Biometeorology* **54**, 255–267.
- CAFFARRA, A. & ECCEL, E. (2011). Projecting the impacts of climate change on the phenology of grapevine in a mountain area. *Australian Journal of Grape and Wine Research* **17**, 52–61.
- CAMPS, J. O. & RAMOS, M. C. (2012). Grape harvest and yield responses to inter-annual changes in temperature and precipitation in an area of north-east Spain with a Mediterranean climate. *International Journal of Biometeorology* **56**, 853–864.
- CHEVET, J. M., LECOCCQ, S. & VISSER, M. (2011). Climate, grapevine phenology, wine production, and prices: Pauillac (1800–2009). *American Economic Review* **101**, 142–146.
- COSTA, A. C., SANTOS, J. A. & PINTO, J. G. (2012). Climate change scenarios for precipitation extremes in Portugal. *Theoretical and Applied Climatology* **108**, 217–234.
- DALLA MARTA, A., GRIFONI, D., MANCINI, M., STORCHI, P., ZIPOLI, G. & ORLANDINI, S. (2010). Analysis of the relationships between climate variability and grapevine phenology in the Nobile di Montepulciano wine production area. *Journal of Agricultural Science, Cambridge* **148**, 657–666.
- DE CORTAZAR-ATAURI, I. G., BRISSON, N. & GAUDILLERE, J. P. (2009). Performance of several models for predicting budburst date of grapevine (*Vitis vinifera* L.). *International Journal of Biometeorology* **53**, 317–326.
- DE ORDUNA, R. M. (2010). Climate change associated effects on grape and wine quality and production. *Food Research International* **43**, 1844–1855.
- DUCHENE, E. & SCHNEIDER, C. (2005). Grapevine and climatic changes: a glance at the situation in Alsace. *Agronomy for Sustainable Development* **25**, 93–99.
- DUCHENE, E., HUARD, F., DUMAS, V., SCHNEIDER, C. & MERDINOGLU, D. (2010). The challenge of adapting grapevine varieties to climate change. *Climate Research* **41**, 193–204.
- FERNANDEZ-GONZALEZ, M., ESCUREDO, O., RODRIGUEZ-RAJO, F. J., AIRA, M. J. & JATO, V. (2011). Prediction of grape production by grapevine cultivar Godello in north-west Spain. *Journal of Agricultural Science, Cambridge* **149**, 725–736.
- FILA, G., GARDIMAN, M., BELVINI, P., MEGGIO, F. & PITACCO, A. (2014). A comparison of different modelling solutions for studying grapevine phenology under present and future climate scenarios. *Agricultural and Forest Meteorology* **195–196**, 192–205.
- FRAGA, H., AMRAOUI, M., MALHEIRO, A. C., MOUTINHO-PEREIRA, J., EIRAS-DIAS, J., SILVESTRE, J. & SANTOS, J. A. (2014a). Examining the relationship between the enhanced

- vegetation index and grapevine phenology. *European Journal of Remote Sensing* **47**, 753–771.
- FRAGA, H., MALHEIRO, A. C., MOUTINHO-PEREIRA, J., CARDOSO, R. M., SOARES, P. M. M., CANCELA, J. J., PINTO, J. G. & SANTOS, J. A. (2014b). Integrated analysis of climate, soil, topography and vegetative growth in Iberian viticultural regions. *Plos ONE* **9**, e108078. DOI: 10.1371/journal.pone.0108078.
- FRAGA, H., MALHEIRO, A. C., MOUTINHO-PEREIRA, J. & SANTOS, J. A. (2014c). Climate factors driving wine production in the Portuguese Minho region. *Agricultural and Forest Meteorology* **185**, 26–36.
- FRAGA, H., COSTA, R., MOUTINHO-PEREIRA, J., CORREIA, C. M., DINIS, L.-T., GONÇALVES, I., SILVESTRE, J., EIRAS-DIAS, J., MALHEIRO, A. C. & SANTOS, J. A. (2015a). Modeling phenology, water status, and yield components of three Portuguese grapevines using the STICS crop model. *American Journal of Enology and Viticulture*. doi: 10.5344/ajev.2015.15031.
- FRAGA, H., SANTOS, J. A., MALHEIRO, A. C., OLIVEIRA, A. A., MOUTINHO-PEREIRA, J. & JONES, G. V. (2015b). Climatic suitability of Portuguese grapevine varieties and climate change adaptation. *International Journal of Climatology*. doi: 10.1002/joc.4325.
- HALL, A. & JONES, G. V. (2009). Effect of potential atmospheric warming on temperature-based indices describing Australian winegrape growing conditions. *Australian Journal of Grape and Wine Research* **15**, 97–119.
- HANNAH, L., ROEHRDANZ, P. R., IKEGAMI, M., SHEPARD, A. V., SHAW, M. R., TABOR, G., ZHI, L., MARQUET, P. A. & HIJMANS, R. J. (2013). Climate change, wine, and conservation. *Proceedings of the National Academy of Sciences of the United States of America* **110**, 6907–6912.
- IPCC (2012). *Managing the risks of extreme events and disasters to advance climate change Adaptation. A Special Report of Working Groups I and II of the Intergovernmental Panel on Climate Change* (Eds C. B. Field, V. Barros, T. F. Stocker, D. Qin, D. J. Dokken, K. L. Ebi, M. D. Mastrandrea, K. J. Mach, G.-K. Plattner, S. K. Allen, M. Tignor, & P. M. Midgley). Cambridge, UK, and New York: Cambridge University Press.
- IVV (2013). *Vinhos e Aguardentes de Portugal, Anuário 2013*. Lisbon, Portugal: Instituto da Vinha e do Vinho.
- JACOB, D., PETERSEN, J., EGGERT, B., ALIAS, A., CHRISTENSEN, O. B., BOUWER, L. M., BRAUN, A., COLETTE, A., DEQUE, M., GEORGIEVSKI, G., GEORGOPOULOU, E., GOBIET, A., MENUT, L., NIKULIN, G., HAENSLER, A., HEMPELMANN, N., JONES, C., KEULER, K., KOVATS, S., KRONER, N., KOTLARSKI, S., KRIEGSMANN, A., MARTIN, E., VAN MEIJAARD, E., MOSELEY, C., PFEIFER, S., PREUSCHMANN, S., RADERMACHER, C., RADTKE, K., RECHID, D., ROUNSEVELL, M., SAMUELSSON, P., SOMOT, S., SOUSSANA, J. F., TEICHMANN, C., VALENTINI, R., VAUTARD, R., WEBER, B. & YIOU, P. (2014). EURO-CORDEX: new high-resolution climate change projections for European impact research. *Regional Environmental Change* **14**, 563–578.
- JONES, G. V. (2005). Climate change in the western United States grape growing regions. *Proceedings of the Seventh International Symposium on Grapevine Physiology and Biotechnology* **689**, 41–59.
- JONES, G. V. & DAVIS, R. E. (2000). Climate influences on grapevine phenology, grape composition, and wine production and quality for Bordeaux, France. *American Journal of Enology and Viticulture* **51**, 249–261.
- JONES, G. V., WHITE, M. A., COOPER, O. R. & STORCHMANN, K.-H. (2004a). Climate and wine: quality issues in a warmer world. In *Proceedings of the Vineyard Data Quantification Society's 11th Enometrics Meeting*. Dijon, France: Vineyard Data Quantification Society. Available online from: http://www.sou.edu/assets/envirostudies/gjones_docs/VDQS%20Climate%20Change.pdf (Verified 25 August 2015).
- JONES, G. V., DUCHÊNE, E., TOMASI, D., YUSTE, J., BRASLAVSKA, O., SCHULTZ, H. R., MARTINEZ, C., BOSO, S., LANGELLIER, F., PERRUCHOT, C. & GUMBERTEAU, G. (2005a). Changes in European winegrape phenology and relationships with climate. In *Proceedings of the XIV GESCO Symposium* (Eds H. Schultz), pp. 54–61. Geisenheim, Germany: Groupe d'Etudes des Systèmes de Conduite de la vigne (GiESCO).
- JONES, G. V., WHITE, M. A., COOPER, O. R. & STORCHMANN, K. (2005b). Climate change and global wine quality. *Climatic Change* **73**, 319–343.
- KELLER, M. (2010a). Managing grapevines to optimise fruit development in a challenging environment: a climate change primer for viticulturists. *Australian Journal of Grape and Wine Research* **16**, 56–69.
- KELLER, M. (2010b). *The Science of Grapevines: Anatomy and Physiology*. Amsterdam: Elsevier.
- KOTLARSKI, S., KEULER, K., CHRISTENSEN, O. B., COLETTE, A., DEQUE, M., GOBIET, A., GOERGEN, K., JACOB, D., LUTHI, D., VAN MEIJAARD, E., NIKULIN, G., SCHAR, C., TEICHMANN, C., VAUTARD, R., WARRACH-SAGI, K. & WULFMEYER, V. (2014). Regional climate modeling on European scales: a joint standard evaluation of the EURO-CORDEX RCM ensemble. *Geoscientific Model Development* **7**, 1297–1333.
- LORENZ, D. H., EICHHORN, K. W., BLEIHOLDER, H., KLOSE, R., MEIER, U. & WEBER, E. (1995). Growth stages of the grapevine: phenological growth stages of the grapevine (*Vitis vinifera* L. ssp. *vinifera*) – codes and descriptions according to the extended BBCH scale. *Australian Journal of Grape and Wine Research* **1**, 100–103.
- MALHEIRO, A. C., SANTOS, J. A., FRAGA, H. & PINTO, J. G. (2010). Climate change scenarios applied to viticultural zoning in Europe. *Climate Research* **43**, 163–177.
- MALHEIRO, A. C., CAMPOS, R., FRAGA, H., EIRAS-DIAS, J. E., SILVESTRE, J. & SANTOS, J. A. (2013). Winegrape phenology and temperature relationships in the Lisbon Wine Region, Portugal. *Journal International des Sciences de la Vigne et du Vin* **47**, 287–299.
- MARTIN, S. R. & DUNN, G. M. (2000). Effect of pruning time and hydrogen cyanamide on budburst and subsequent phenology of *Vitis vinifera* L. variety Cabernet Sauvignon in central Victoria. *Australian Journal of Grape and Wine Research* **6**, 31–39.
- MCINTYRE, G. N., LIDER, L. A. & FERRARI, N. L. (1982). The chronological classification of grapevine phenology. *American Journal of Enology and Viticulture* **33**, 80–85.
- MORIONDO, M. & BINDI, M. (2007). Impact of climate change on the phenology of typical mediterranean crops. *Italian Journal of Agrometeorology* **2007**, 5–12.

- MORIONDO, M., JONES, G. V., BOIS, B., DIBARI, C., FERRISE, R., TROMBI, G. & BINDI, M. (2013). Projected shifts of wine regions in response to climate change. *Climatic Change* **119**, 825–839.
- MORIONDO, M., FERRISE, R., TROMBI, G., BRILLI, L., DIBARI, C. & BINDI, M. (2015). Modelling olive trees and grapevines in a changing climate. *Environmental Modelling & Software*. doi: 10.1016/j.envsoft.2014.12.016.
- NENDEL, C. (2010). Grapevine bud break prediction for cool winter climates. *International Journal of Biometeorology* **54**, 231–241.
- NEUMANN, P. A. & MATZARAKIS, A. (2014). Estimation of wine characteristics using a modified Heliothermal Index in Baden-Württemberg, SW Germany. *International Journal of Biometeorology* **58**, 407–415.
- OIV (2009). *Code des Caractères Descriptifs des Variétés et Espèces de Vitis*, 2nd éd. Paris, France: Organisation Internationale de la Vigne et du Vin.
- OLSSON, C. & JÖNSSON, A. M. (2015). Budburst model performance: the effect of the spatial resolution of temperature data sets. *Agricultural and Forest Meteorology* **200**, 302–312.
- PARKER, A. K., DE CORTAZAR-ATAURI, I. G., VAN LEEUWEN, C. & CHUINE, I. (2011). General phenological model to characterise the timing of flowering and veraison of *Vitis vinifera* L. *Australian Journal of Grape and Wine Research* **17**, 206–216.
- PARKER, A., DE CORTAZAR-ATAURI, I. G., CHUINE, I., BARBEAU, G., BOIS, B., BOURSQUOT, J.-M., CAHUREL, J.-Y., CLAVERIE, M., DUFOURCQ, T., GÉNY, L., GUIMBERTEAU, G., HOFMANN, R. W., JACQUET, O., LACOMBE, T., MONAMY, C., OJEDA, H., PANIGAL, L., PAYAN, J.-C., LOVELLE, B. R., ROUCHAUD, E., SCHNEIDER, C., SPRING, J.-L., STORCHI, P., TOMASI, D., TRAMBOUZE, W., TROUGHT, M. & VAN LEEUWEN, C. (2013). Classification of varieties for their timing of flowering and veraison using a modelling approach: a case study for the grapevine species *Vitis vinifera* L. *Agricultural and Forest Meteorology* **180**, 249–264.
- PEEL, M. C., FINLAYSON, B. L. & MCMAHON, T. A. (2007). Updated world map of the Köppen-Geiger climate classification. *Hydrology and Earth System Sciences* **11**, 1633–1644.
- PETRIE, P. R. & SADRAS, V. O. (2008). Advancement of grapevine maturity in Australia between 1993 and 2006: putative causes, magnitude of trends and viticultural consequences. *Australian Journal of Grape and Wine Research* **14**, 33–45.
- REAL, A. C., BORGES, J., CABRAL, J. S. & JONES, G. V. (2015). Partitioning the grapevine growing season in the Douro Valley of Portugal: accumulated heat better than calendar dates. *International Journal of Biometeorology* **59**, 1045–1059.
- SADRAS, V. O. & MORAN, M. A. (2013). Nonlinear effects of elevated temperature on grapevine phenology. *Agricultural and Forest Meteorology* **173**, 107–115.
- SANTOS, J. & CORTE-REAL, J. (2006). Temperature extremes in Europe and wintertime large-scale atmospheric circulation: HadCM3 future scenarios. *Climate Research* **31**, 3–18.
- SANTOS, J. A., CORTE-REAL, J., ULBRICH, U. & PALUTIKOF, J. (2007). European winter precipitation extremes and large-scale circulation: a coupled model and its scenarios. *Theoretical and Applied Climatology* **87**, 85–102.
- SANTOS, J. A., GRÄTSCH, S. D., KARREMANN, M. K., JONES, G. V. & PINTO, J. G. (2013). Ensemble projections for wine production in the Douro Valley of Portugal. *Climatic Change* **117**, 211–225.
- TOMASI, D., JONES, G. V., GIUSTI, M., LOVAT, L. & GAIOTTI, F. (2011). Grapevine Phenology and climate change: relationships and trends in the Veneto Region of Italy for 1964–2009. *American Journal of Enology and Viticulture* **62**, 329–339.
- VAN LEEUWEN, C., GARNIER, C., AGUT, C., BACULAT, B., BARBEAU, G., BESNARD, E., BOIS, B., BOURSQUOT, J.-M., CHUINE, I., DESSUP, T., DUFOURCQ, T., GARCIA-CORTAZAR, I., MARGUERIT, E., MONAMY, C., KOUNDOURAS, S., PAYAN, J.-C., PARKER, A., RENOUF, V., RODRIGUEZ-LOVELLE, B., ROBY, J.-P., TONETTO, J. & TRAMBOUZE, W. (2008). Heat requirements for grapevine varieties is essential information to adapt plant material in a changing climate. In *Proceedings of the VIIème Congrès International des Terroirs Viticoles* (Eds F. Murisier), pp. 222–227. Nyon, Switzerland: Agroscope Changins-Wädenswil ACW.
- WEBB, L. B., WHETTON, P. H. & BARLOW, E. W. R. (2007). Modelled impact of future climate change on the phenology of winegrapes in Australia. *Australian Journal of Grape and Wine Research* **13**, 165–175.
- WEBB, L. B., WHETTON, P. H. & BARLOW, E. W. R. (2008). Modelling the relationship between climate, winegrape price and winegrape quality in Australia. *Climate Research* **36**, 89–98.
- WEBB, L. B., WHETTON, P. H. & BARLOW, E. W. R. (2011). Observed trends in winegrape maturity in Australia. *Global Change Biology* **17**, 2707–2719.
- WEBB, L. B., WHETTON, P. H., BHEND, J., DARBYSHIRE, R., BRIGGS, P. R. & BARLOW, E. W. R. (2012). Earlier winegrape ripening driven by climatic warming and drying and management practices. *Nature Climate Change* **2**, 259–264.
- WHITE, M. A., DIFFENBAUGH, N. S., JONES, G. V., PAL, J. S. & GIORGI, F. (2006). Extreme heat reduces and shifts United States premium wine production in the 21st century. *Proceedings of the National Academy of Sciences of the United States of America* **103**, 11217–11222.
- WILKS, D. S. (2011). *Statistical Methods in the Atmospheric Sciences*. Amsterdam: Academic Press.
- WINKLER, A. J., COOK, J. A., KLIEWER, W. M. & LIDER, L. A. (1974). *General Viticulture*. California: University of California Press.
- ZAPATA, D., SALAZAR, M., CHAVES, B., KELLER, M. & HOOGENBOOM, G. (2015). Estimation of the base temperature and growth phase duration in terms of thermal time for four grapevine cultivars. *International Journal of Biometeorology*. DOI: 10.1007/s00484-015-0985-y.